

## CLAIMS

We claim:

- 5 ~~Sub E~~
- 10
- 15 ~~Sub E~~
- 20
1. A method for the recombination of nucleic acid constructs, comprising:
    - a) providing:
      - i) a first nucleic acid construct comprising, in operable order, an origin of replication, a first sequence-specific recombinase target site, and a nucleic acid of interest;
      - ii) a second nucleic acid construct comprising, in operable order, an origin of replication, a regulatory element and a second sequence-specific recombinase target site adjacent to and downstream from said regulatory element; and
      - iii) a site-specific recombinase;
    - b) contacting said first and said second nucleic acid constructs with said site-specific recombinase under conditions such that said first and second nucleic acid constructs are recombined to form a third nucleic acid construct, wherein said nucleic acid of interest is operably linked to said regulatory element.
  2. The method of Claim 1, wherein said regulatory element comprises a promoter element.
  3. The method of Claim 1, wherein said regulatory element comprises a fusion peptide.
  4. The method of Claim 3, wherein said fusion peptide comprises an affinity domain.

Sub  
E.F.  
cont.

5. The method of Claim 3, wherein said fusion peptide comprises an epitope tag.

6. The method of Claim 1, wherein said nucleic acid of interest comprises a gene.

5 7. The method of Claim 1, wherein said first nucleic acid construct further comprises a selectable marker.

8. The method of Claim 1, wherein said second nucleic acid construct further comprises a selectable marker.

10 9. The method of Claim 1, wherein said first nucleic acid construct further comprises a prokaryotic termination sequence.

10. The method of Claim 1, wherein said first nucleic acid construct further comprises a eukaryotic polyadenylation sequence.

11. The method of Claim 1, wherein said first nucleic acid construct further comprises a conditional origin of replication.

15 12. The method of Claim 1, wherein said first sequence-specific recombinase target site is selected from the group consisting of *loxP*, *loxP2*, *loxP3*, *loxP23*, *loxP511*, *loxB*, *loxC2*, *loxL*, *loxR*, *loxΔ86*, *loxΔ117*, *frt*, *dif*, *loxH* and *att*.

20 13. The method of Claim 1, wherein said second sequence-specific recombinase target site is selected from the group consisting of *loxP*, *loxP2*, *loxP3*, *loxP23*, *loxP511*, *loxB*, *loxC2*, *loxL*, *loxR*, *loxΔ86*, *loxΔ117*, *frt*, *dif*, *loxH* and *att*.

14. The method of Claim 1, wherein said first nucleic acid construct further comprises a polylinker.

15. The method of Claim 1, wherein said contacting said first and said second nucleic acid constructs with said site-specific recombinase comprises introducing said first and said second nucleic acid constructs into a host cell under conditions such that said third nucleic acid construct is capable of replicating in said host cell.

16. The method of Claim 15, wherein said site-specific recombinase is encoded by said host cell.

17. The method of Claim 1, wherein said first nucleic acid construct further comprises a third sequence-specific recombinase target site and said second nucleic acid constructs further comprises a fourth sequence-specific recombinase target site.

18. The method of Claim 17, wherein said first sequence-specific recombinase target site and said third sequence-specific recombinase target site in said first nucleic acid construct are located on opposite sides of said nucleic acid of interest.

19. The method of Claim 17, wherein in said third and fourth sequence-specific recombinase target sites are selected from the group consisting of RS sites and Res sites.

20. The method of Claim 1, wherein said first nucleic acid construct further comprises a third sequence-specific recombinase target site and said second nucleic acid constructs further comprises a fourth sequence-specific recombinase target site, wherein the method further comprises providing a second site-specific recombinase and step c) contacting said third nucleic acid construct with said second site-specific

recombinase under conditions such that said third nucleic acid construct is recombined to form a fourth and a fifth nucleic acid construct.

21. A recombined nucleic acid construct prepared according to the method of Claim 1.

22. A method for the recombination of nucleic acid constructs, comprising:

a) providing:

i) a vector;

ii) a linear nucleic acid molecule comprising a sequence complementary to at least a portion of said vector; and

iii) an *E. coli* host cell, wherein said host cell comprises an endogenous recombination system, a loss of function *rec* mutation, a suppressor, and a loss of function endogenous restriction modification system mutation; and

b) introducing said vector and said linear nucleic acid molecule into said host cell under conditions such that said linear nucleic acid molecule and said vector are recombined to form a recombinant nucleic acid construct.

23. The method of Claim 22, wherein said loss of function *rec* mutation is selected from the group consisting of *recBC* and *recD*.

24. The method of Claim 22, wherein said suppressor comprises *sbc*.

25. The method of Claim 22, wherein said loss of function endogenous restriction modification system mutation comprises *hsdR*.

26. A method for the cloning of nucleic acid libraries, comprising:

a) providing:

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995

i) a plurality of first nucleic acid constructs comprising, in operable order, an origin of replication, a first sequence-specific recombinase target site, and a nucleic acid member from a nucleic acid library;

ii) a plurality of second nucleic acid constructs comprising, in operable order, an origin of replication, a regulatory element and a second sequence-specific recombinase target site adjacent to and downstream from said regulatory element; and

iii) a site-specific recombinase;

b) contacting said plurality of first and second nucleic acid constructs with said site-specific recombinase under conditions such that said plurality of first and second nucleic acid constructs are recombined to form a plurality of third nucleic acid constructs, wherein said nucleic acid members from said nucleic acid library are operably linked to said regulatory elements.

27. A nucleic acid library prepared according to the method of Claim 26.

28. A method for the directional cloning of a nucleic acid molecule, comprising:

a) providing:

i) first and second portions of a regulatory element;

ii) a first nucleic acid molecule comprising said first portion of said regulatory element; and

iii) a second nucleic acid molecule comprising said second portion of said regulatory element; and

b) combining said first and said second nucleic acid molecules to produce a third nucleic acid molecule under conditions whereby an intact regulatory element is produced from the combination of said first and said second portions of said regulatory element, wherein the presence of said intact regulatory element in said third nucleic acid molecule indicates a direction of

cloning of said first nucleic acid molecule with respect to said second nucleic acid molecule.

29. The method of Claim 28, wherein said regulatory element comprises a *lacO* site.

5                    22  
30. A method for regulated recombination in host cells that constitutively express a recombinase, comprising:

a) providing:

i) a host cell expressing a recombinase;

10                    ii) a first nucleic acid construct comprising an origin of replication, a first site-specific recombinase site, a second site-specific recombinase site that differs in sequence from said first site-specific recombinase site such that said recombinase will not initiate recombination between said first and second site-specific recombinase sites, and a selectable marker gene between said first and second site-specific recombinase sites; and

15                    iii) a second nucleic acid construct comprising an origin of replication, a third site-specific recombinase target site, and a fourth site-specific recombinase target site that differs in sequence from said third site-specific recombinase target site such that said recombinase will not initiate recombination between said third and fourth site-specific recombinase sites; and

20                    b) introducing said first and second nucleic acid constructs into said host cell under conditions such that said first and second nucleic acid constructs are recombined.

25                    23                    22  
31. The method of Claim 30, further comprising the step of selecting for a desired recombinant nucleic acid molecule using said selectable marker.

24  
32. The method of Claim 30<sup>22</sup>, wherein said first nucleic acid construct is a Univector.

26  
33. The method of Claim 30<sup>22</sup>, wherein said second nucleic acid construct is a Univector.

5 26  
34. A host cell expressing a recombinant nucleic acid construct prepared according to the method of Claim 30<sup>22</sup>, wherein said host cell constitutively expresses a recombinase.

27  
35. A method for the recombination of nucleic acid constructs, comprising:

a) providing:

i) a first nucleic acid construct comprising a *loxH* site;

ii) a second nucleic acid construct comprising a *loxH* site;

and

iii) a site-specific recombinase; and

b) contacting said first and said second nucleic acid constructs with said site-specific recombinase under conditions such that said first and second nucleic acid constructs are recombined.

36. A recombined nucleic acid construct prepared according to the method of Claim 35.

ADD E3

ADD G'